arXiv:1110.4026v1 [math.OA] 18 Oct 2011

APERIODICITY CONDITIONS IN TOPOLOGICAL k-GRAPHS

SARAH WRIGHT

ABSTRACT. We give two new conditions on topological k-graphs that are equivalent to the Yeend's aperiodicity Condition (A). Each of the new conditions concerns finite paths rather than infinite. We use a specific example, resulting from a new construction of a twisted topological k-graph, to demonstrate the improvements achieved by the new conditions. Reducing this proof of equivalence to the discrete case also gives a new direct proof of the corresponding conditions in discrete k-graphs, where previous proofs depended on simplicity of the corresponding C*-algebra.

1. INTRODUCTION

The theory of graph C*-algebras began with the work of Cuntz and Kreiger [1] and the later work of Enomoto and Watatani [2]. Since then, there have been many contributions by a variety of researchers resulting in an extensive collection of literature. The main idea is to associate to a directed graph, a C*-algebra and use the combinatorics of the graph to answer questions on the structure of the C*-algebra. The study of graph algebras had been particularly fruitful in its provision of a rich class of easily accessible examples of C*-algebras with various properties.

To this end, the idea of a graph has been generalized in a few ways including the k-graphs of Kumjian and Pask [5] and the topological graphs of Katsura [4]. Most recently, the work of Yeend provides a generalization of both higher-rank graphs and topological graphs, with the unifying theory of topological k-graphs [11].

An important outcome of the study of each type of graph is the relationship between the periodic paths (or lack there of) in the graph and the simplicity of the associated C*-algebra. The first such aperiodicity condition for directed graphs appeared in [6], there referred to as Condition (L), and states that every cycle of the graph must have an entry. This is one of the necessary conditions for simplicity and that relationship demonstrates the beauty of the subject; it is easy to look at picture of a directed graph and check if each cycle has an entry.

For the best viewing of the figures, it is recommended that the reader print the document in color.

The key to cycles having or not having entries is the idea of being able to "back out of a cycle" and build an aperiodic infinite path. As the infinite paths play such an important role in the groupoid construction of the C*-algebra, in the case of higher-rank graphs the original aperiodicity condition, called Condition (A), was stated in terms of infinite paths. Lewin, Robertson, and Sims have developed conditions on higher-rank graphs, which are equivalent to aperiodicity [8, 7], that have proven easier to work with in examples and in general proofs. The work of this paper gives extensions of the various aperiodicity conditions to the case of topological k-graphs and proves their equivalence.

In the first section we'll discuss the background on topological k-graphs that is needed. In the middle section we state the two new aperiodicity conditions and prove the main result of equivalence with Yeend's Condition (A). We also address a new proof of the equivalence of the corresponding discrete conditions. The final section gives a new method of constructing a topological k-graph from a discrete k-graph. The construction of these twisted topological k-graphs shares the flavor of Yeend's skew product graphs [11, Definition 8.1] as well as the topological dynamical systems defined by Farthing, Patani, and Willis, [3]. Some advantages include the fact that we need not begin with a topological k-graph, but can use a discrete graph with desired properties. Also, there are fewer requirements of the spaces and maps involved, making it easier to construct useful examples. We give a specific example of this construction and use one of the new conditions to determine the topological k-graph is aperiodic, demonstrating the improvements gained by considering finite paths.

Acknowledgement: I would like to give special thanks to Aidan Sims for many helpful conversations, ideas, and motivation.

2. Background

The basics (and considerably more) on topological k-graphs and their C*-algebras can be found in Yeend's original papers [12, 11, 10]. We review some of the basics here for convenience.

We will regard $\mathbb{N}^k$ as the category with a single object and composition given by addition. We use $\{e_i\}_{i=1}^k$ to represent the standard basis of $\mathbb{N}^k$ and for $m \in \mathbb{N}^k$ denote the i^{th} component by m_i. For $m, n \in \mathbb{N}^k$, we say $m \geq n$ if $m_i \geq n_i$ for every $i \in \{1, 2, \ldots, k\}$, write $m \vee n$ for the coordinate maximum, and $m \wedge n$ for the coordinate minimum.

For a natural number k, a *topological k-graph* is a pair (Λ, d) consisting of a small category $\Lambda = (\mathrm{Obj}(\Lambda), \mathrm{Mor}(\Lambda), r, s)$ and a functor $d : \Lambda \to \mathbb{N}^k$ which satisfy the following:

1. $\mathrm{Obj}(\Lambda)$ and $\mathrm{Mor}(\Lambda)$ are second-countable, locally compact, Hausdorff spaces;
2. $r, s : \mathrm{Mor}(\Lambda) \to \mathrm{Obj}(\Lambda)$ are continuous and s is a local homeomorphism;

(3) Composition $\circ : \Lambda \times_c \Lambda \to \Lambda$ is continuous and open, where $\Lambda \times_c \Lambda$ has the relative product topology inherited from the product topology on $\Lambda \times \Lambda$;
(4) $d : \Lambda \to \mathbb{N}^k$ is continuous, where $\mathbb{N}^k$ has the discrete topology;
(5) For all $\lambda \in \Lambda$ and all $m, n \in \mathbb{N}^k$ such that $d(\lambda) = m + n$, there exist unique $\mu, \nu \in \Lambda$ such that $d(\mu) = m$, $d(\nu) = n$, and $\lambda = \mu\nu$.

Condition (5) is called the *unique factorization property* and is exactly the same as that for discrete k-graphs. The functor d is called the *degree map*, or the *shape map*. For $n \in \mathbb{N}^k$, we denote by Λ^n the set $d^{-1}(n)$ in $\mathrm{Mor}(\Lambda)$, and refer to the elements as *paths of shape n*. It is often necessary to deal with the range and source maps as they relate to paths of a specific shape, so the notation r_n refers to the restriction of the range map $r_n|_{\Lambda^n}$ and similarly $s_n := s|_{\Lambda^n}$. A consequence of the unique factorization property is that Λ^0 is the same as the identity morphisms of Λ, and these are thus referred to as the *vertices* of Λ. For any sets $X, Y \subset \Lambda$ we use XY for the set $\{\mu\nu \mid \mu \in X, \nu \in Y, s(\mu) = r(\nu)\}$. (Notice that we concatenate paths so that the range of the later path equals the source of the previous path. This may seem counter-intuitive to those new to subject, but it is considered the standard notation in higher-rank graphs, and makes more sense when constructing the C*-algebra.) This notation is of particular use when the first set is a set of vertices and the second set is all the paths. So, for $V \subset \Lambda^0$, $V\Lambda = \{\lambda \in \Lambda \mid r(\lambda) \in V\}$ and $\Lambda V = \{\lambda \in \Lambda \mid s(\lambda) \in V\}$.

The factorization property implies that for $\lambda \in \Lambda$ and $m < n < d(\lambda) \in \mathbb{N}^k$ there are unique paths denoted $\lambda(0, m) \in \Lambda^m$, $\lambda(m, n) \in \Lambda^{n-m}$, and $\lambda(n, d(\lambda)) \in \Lambda^{d(\lambda)-n}$ such that $\lambda(0, m)\lambda(m, n)\lambda(n, d(\lambda)) = \lambda$. We therefore think of $\lambda(p, q)$ as the portion of the path λ that runs from q to p.

An important class of examples of k-graphs are the *grid graphs*, $\Omega_{k,m}$. For a fixed $k \geq 1$ and $m \in (\mathbb{N} \cup \{\infty\})^k$ the discrete k-graph $\Omega_{k,m}$ has morphisms $\{(p, q) \in \mathbb{N}^k \times \mathbb{N}^k \mid p \leq q \leq m\}$, range and source maps given by $r(p, q) = (p, p)$ and $s(p, q) = (q, q)$, composition defined as $(p, q)(q, r) = (p, r)$ and degree map given by $d(p, q) = q - p$.

We visualize discrete k-graphs by their *1-skeletons*, a directed graph whose vertices are Λ^0 and whose edges are paths from the sets Λ^{e_i} and are colored k different colors depending on the shape. In some cases, such as any $\Omega_{k,m}$, we can construct the entire k-graph from the 1-skeleton. In other cases we need more information. Consider the path bd in Figure 1(b) of shape $(0, 1, 1)$. By the unique factorization property, bd must factor uniquely as a "red-blue" path. So, $bd = \mu\nu$ with $d(\mu) = (0, 0, 1)$ and $d(\nu) = (0, 1, 0)$. It cannot be determined from the 1-skeleton alone if the unique factorization we desire is



(a) The 1-Skeleton of $\Omega_{3,(3,2,1)}$

(b) A 1-skeleton of a 3-graph requiring factorization rules

Figure 1. 1-Skeletons of two different 3-graphs

cb or cb'. We need to specify factorization rules for each bi-colored path:

$$\begin{aligned} da &= a'c \\ fa &= a'e \\ bd &= cb' \\ bf &= eb' \\ fd &= df \\ ce &= ec. \end{aligned}$$

For graphs of rank k, (Λ_1, d_1) and (Λ_2, d_2), a *topological k-graph morphism* is a continuous degree preserving functor $x : \Lambda_1 \to \Lambda_2$. That is, $d_2(x(\lambda)) = d_1(\lambda)$. We are particularly concerned with the graph morphisms from the grid graphs. The *path space* of a topological k-graph Λ

$$X_\Lambda := \bigcup_{m \in (\mathbb{N} \cup \{\infty\})^k} \{x : \Omega_{k,m} \to \Lambda \,|\, x \text{ is a graph morphism}\}$$

can be thought to include Λ since any finite path λ uniquely determines a graph morphism $x_\lambda : \Omega_{k,d(\lambda)} \to \Lambda$ such that $x(m,n) = \lambda(m,n)$ for any $m \le n \le d(\lambda)$. We extend the idea of range and degree so that $r(x) = x(0,0)$ and $d(x) = m$ for $x : \Omega_{k,m} \to \Lambda$. In practice, we think of taking a particular grid graph and labeling the vertices and edges while following the factorization rules associated to the 1-skeleton of Λ. So, a path of shape (∞, ∞, ∞) with range v in the 3-graph of Figure 1(b) with the factorization rules given could look like that of Figure 2.

In order to define aperiodic paths in a topological k-graph we need the idea of the shift map. The *shift map* σ^m on X_Λ assigns to x the unique path $\sigma^m x$ in X_Λ which satisfies $d(\sigma^m x) = d(x) - m$ and $\sigma^m x(0,n) = x(m, m+n)$ for all $0 \leqslant n \leqslant d(x) - m$. For $m \in \mathbb{N}^k$, the shift map of degree m removes a path of shape m from the range end of x. We say a path $x \in X_\Lambda$ is *aperiodic* if for any $m, n \in \mathbb{N}^k$ $\sigma^m x = \sigma^n x$ implies that $m = n$.

The idea of a minimal common extension of two paths will be important in one of our formulations of aperiodicity, but it is also necessary in our



Figure 2. An Infinite Path in a 3-Graph

definition of a boundary path. For paths $\mu, \nu \in \Lambda$, we say λ is a *common extension of* μ and ν if we can factor $\lambda = \mu\mu'$ and $\lambda = \nu\nu'$ for some $\mu', \nu' \in \Lambda$. We consider λ to be a *minimal common extension* if it also satisfies $d(\lambda) = d(\mu) \vee d(\nu)$. We denote by $\mathrm{MCE}(\mu, \nu)$ the set of minimal common extensions of μ and ν and for subsets $X, Y \subset \Lambda$, and define

$$\mathrm{MCE}(X, Y) := \bigcup_{\mu \in X, \nu \in Y} \mathrm{MCE}(\mu, \nu).$$

For a vertex $v \in \Lambda^0$, we say a subset $E \subset \Lambda$ is *compact exhaustive* for v if E is compact, $r(E)$ is a neighborhood of v, and for all $\lambda \in r(E)\Lambda$ there exists a $\mu \in E$ such that $\mathrm{MCE}(\lambda, \mu) \neq \emptyset$. A path $x \in X_\Lambda$ is considered a *boundary path* if for any $m \in \mathbb{N}^k$ with $m \leq d(x)$ and any set E which is compact exhaustive set for $x(m, m)$, there exists a $\lambda \in E$ such that $x(m, m + d(\lambda)) = \lambda$. We write $\partial\Lambda$ for the set of all boundary paths in X_Λ. As in discrete k-graphs the boundary paths are a generalization of the infinite paths of directed graphs, [9, Lemma 4.22].

Under the condition that Λ is *compactly aligned*, that is if both $X, Y \subset \Lambda$ are compact, then $\mathrm{MCE}(X, Y)$ is also compact, we can build a topological groupoid $\mathcal{G}_\Lambda$ called the *boundary path groupoid* which has $\partial\Lambda$ as its unit space and can be endowed with a locally compact Hausdorff topology. The C*-algebra of Λ, $C^*(\Lambda)$, is $C^*(\mathcal{G}_\Lambda)$, the full groupoid C*-algebra of the boundary path groupoid. As we will not need the details of the construction of the C*-algebra to understand the results of this paper, we refer the reader to [11] for the formulation of this topology as well as a more in depth discussion. However, to demonstrate the importance of aperiodicity, we conclude the background information with a result of Yeend.

Proposition 2.1. [11, Theorem 5.2] *For a compactly aligned topological k-graph (Λ, d), the boundary path groupoid $\mathcal{G}_\Lambda$ is topologically principal if and only if for every open set $V \subset \Lambda^0$ there exists an $x \in V\partial\Lambda$ which is aperiodic.*

3. Results

The condition that every open set of vertices must have an aperiodic path that terminates in that set is the natural extension of aperiodicity conditions for topological graphs and discrete k-graphs. Yeend introduced the condition and referred to it as Condition (A). Our main result, Theorem 3.1, gives two new equivalent conditions to Yeend's Condition (A).

Theorem 3.1. *Let (Λ, d) be a compactly aligned topological k-graph and V a nonempty open subset of Λ^0. The following conditions are equivalent.*

(A) There exists an aperiodic path $x \in V\partial\Lambda$.

(B) For any pair $m \neq n \in \mathbb{N}^k$ there exists a path $\lambda_{V,m,n} \in V\Lambda$ such that $d(\lambda) \geq m \vee n$ and

$$(\star) \qquad \lambda(m, m + d(\lambda) - (m \vee n)) \neq \lambda(n, n + d(\lambda) - (m \vee n)).$$

(C) There is a vertex $v \in V$ and paths $\alpha, \beta \in \Lambda$ with $s(\alpha) = s(\beta) = v$ such that there exists a path $\tau \in s(\alpha)\Lambda$ with $\mathrm{MCE}(\alpha\tau, \beta\tau) = \emptyset$.

It is not difficult to see that in the discrete case, (if $\mathrm{Obj}(\Lambda)$ and $\mathrm{Mor}(\Lambda)$ both have the discrete topology) then Conditions (A) – (C) above reduce to [5, Definition 4.3], Condition (iv) of [8, Lemma 3.2], and [7, Definition 3.1] respectively. This special case gives the following corollary. The equivalence that appears in Corollary 3.2 is already known, as each condition is a necessary condition for simplicity of the associated C*-algebra. The direct proof of this equivalence that follows from the proof of Theorem 3.1 is new, and an important outcome of this work.

Corollary 3.2. *Let (Λ, d) be a discrete finitely aligned k-graph. Then the following are equivalent.*

(i) For each $v \in \Lambda^0$ there exists and aperiodic path $x \in v\Lambda^\infty$.

(ii) For each $v \in \Lambda^0$ and each $m \neq n \in \mathbb{N}^k$ there exists a path $\lambda \in v\Lambda$ such that $d(\lambda) \geq m \vee n$ and which satisfies $(\star)$.

(iii) For every pair of distinct paths $\alpha, \beta \in \Lambda$ with $s(\alpha) = s(\beta)$ there exists a path $\tau \in s(\alpha)\Lambda$ such that $\mathrm{MCE}(\alpha\tau, \beta\tau) = \emptyset$.

The proof of the theorem requires an important lemma. Lemma 3.3 is an extension of Lemma 5.6 of [4] which shows that near a path which satisfies $(\star)$ are a lot of other paths which also satisfy $(\star)$. This allows us to extend the proof of the equivalence of (i) and (iv) in Lemma 3.2 of [8] to topological k-graphs.

Lemma 3.3. *Let V be a nonempty open subset of Λ^0, $m \neq n \in \mathbb{N}^k$, and $\lambda \in V\Lambda$. If λ satisfies $(\star)$ for m and n, then there exists a compact neighborhood $E \subset V\Lambda^{d(\lambda)}$ of λ such that every $\mu \in E$ satisfies $(\star)$.*

Proof. Suppose $\lambda \in V\Lambda$ satisfies $(\star)$. We can factor λ in two ways:

$$\lambda = \lambda(0, m)\lambda(m, m + d(\lambda) - (m \vee n))\lambda(m + d(\lambda) - (m \vee n), d(\lambda)) \text{ and}$$
$$\lambda = \lambda(0, n)\lambda(n, n + d(\lambda) - (m \vee n))\lambda(n + d(\lambda) - (m \vee n)), d(\lambda))$$

Let E_m and E_n be disjoint compact neighborhoods of $\lambda(m, m + d(\lambda) - (m \vee n))$ and $\lambda(n, n + d(\lambda) - (m \vee n))$. Also, let E_1, E_2, E_3, and E_4 be compact neighborhoods of $\lambda(0, m)$, $\lambda(m + d(\lambda) - (m \vee n), d(\lambda))$, $\lambda(0, n)$, and $\lambda(n + d(\lambda) - (m \vee n), d(\lambda))$ respectively. We define the set E' to be the paths in $\Lambda^{d(\lambda)}$ that can be factored as paths in E_1, E_m and E_2 as well as E_3, E_n, and E_4. So,

$$E' := \{\alpha\beta\gamma : (\alpha, \beta, \gamma) \in E_1 \times_c E_m \times_c E_2\} \cap \{\mu\nu\xi : (\mu, \nu, \xi) \in E_3 \times_c E_n \times_c E_4\} \subset \Lambda^{d(\lambda)}.$$

Now, let $F \subset V$ be a compact neighborhood of $r(\lambda)$ and

$$E = E' \cap r^{-1}_{d(\lambda)}(F).$$

Then $E \subset V\Lambda^{d(\lambda)}$ is a compact neighborhood of λ with nonempty interior in which every element satisfies $(\star)$. □

Proof of Theorem 3.1. We'll show Condition (B) is equivalent to each of (A) and (C).

(A) $\implies$ (B). Fix and open set $V \subset \Lambda^0$, and a pair $m \neq n \in \mathbb{N}^k$, and suppose $x \in \partial\Lambda$ is aperiodic. Then, $\sigma^m x \neq \sigma^n x$ and so for a sufficiently large $p \in \mathbb{N}^k$

$$\sigma^m x(0, p) \neq \sigma^n x(0, p).$$

Let $\lambda = x(0, p + m \vee n)$, then

$$\begin{aligned} \lambda(m, m + d(\lambda) - m \vee n) &= x(m, m + (p + m \vee n) - m \vee n) \\ &= \sigma^m x(0, p) \\ &\neq \sigma^n x(0, p) \\ &= x(n, n + p + m \vee n - m \vee n) \\ &= \lambda(n, n + d(\lambda) - m \vee n). \end{aligned}$$

So, $d(\lambda) \geq m \vee n$ and λ satisfies $(\star)$.

(B) $\implies$ (A). Fix an open set $V \subseteq \Lambda^0$ and let $\{(m_i, n_i)\}_{i=1}^{\infty}$ be a listing of the elements in the set $\{(m, n) \in \mathbb{N}^k \times \mathbb{N}^k : m \neq n\}$. Let $V_1 = V$, choose $\lambda_1 \in V_1\Lambda$ such that $d(\lambda_1) \geq m_1 \vee n_1$ and $(\star)$ is satisfied for m_1 and n_1; choose a compact neighborhood F_1 of λ_1 by Lemma 3.3; and let $E_1 := F_1\partial\Lambda = \{x \in \partial\Lambda : x\,(0, d(\lambda_1)) \in F_1\}$. Now, proceed inductively:

$V_i :=$ interior of $s(F_{i-1})$,

$\lambda_i := \lambda_{V_i, m_i, n_i}$ satisfy $(\star)$ for m_i and n_i,

$F_i :=$ a compact neighborhood of λ_i given by Lemma 3.3, and

$E_i := F_1 \dots F_i\partial\Lambda$.

Notice that each E_i is compact in $\partial\Lambda$ by Lemma 3.8 of [10]: Let $p = q = \sum_{j=1}^{i} d(\lambda_j)$ and $U = V = F_1 \dots F_i \subset \Lambda^{\sum_{j=1}^{i} d(\lambda_j)}$. Then E_i is exactly the set $\mathcal{Z}(U *_s V, p-q)$, which is compact in $\partial\Lambda$. Also note that the E_i's are nested; $E_1 \supset E_2 \supset \ldots$. Now, let

$$E := \bigcap_{i=1}^{\infty} E_i \subset \partial\Lambda.$$

The set E is non-empty, so take any $x \in E$. Let $\mu_i = x\,(d(\lambda_1\lambda_2\ldots\lambda_{i-1}), d(\lambda_i))$ for each $i \in \mathbb{N}$. Then by Lemma 4.1.3 of [12], x is the unique element in X_Λ such that $d(x) = \lim_{j\to\infty} d(\mu_1\mu_2\ldots\mu_j)$ and $x(0, d(\mu_1\mu_2\ldots\mu_j)) = \mu_1\mu_2\ldots\mu_j$, so it makes sense to write

$$x = \mu_1\mu_2\mu_3\ldots,$$

with each $\mu_i \in F_i$. Fix $m \neq n \in \mathbb{N}^k$. Then for some $i \in \mathbb{N}$, $(m,n) = (m_i, n_i)$ in the listing above. So,

$$\begin{aligned}\sigma^m x\left(\sum_{j=1}^{i-1} d(\lambda_j), \sum_{j=1}^{i} d(\lambda_j) - (m \vee n)\right) &= \sigma^{m+\sum_{j=1}^{i-1} d(\lambda_j)} x(0, d(\lambda_i) - (m \vee n)) \\ &= \sigma^{\sum_{j=1}^{i-1} d(\lambda_j)} x(m, m + d(\lambda_i) - (m \vee n)) \\ &= \mu_i(m, m + d(\mu_i) - (m \vee n)).\end{aligned}$$

Similarly,

$$\sigma^n x\left(\sum_{j=1}^{i-1} d(\lambda_j), \sum_{j=1}^{i} d(\lambda_j) - (m \vee n)\right) = \mu_i(n, n + d(\mu_i) - (m \vee n)).$$

By the definition of F_i, we know that

$$\mu_i(m, m + d(\mu_i) - (m \vee n)) \neq \mu_i(n, n + d(\mu_i) - (m \vee n)),$$

and thus $\sigma^m x \neq \sigma^n x$ and x is an aperiodic boundary path.

(B) $\Longrightarrow$ (C). Fix an open set $V \subset \Lambda^0$ and paths $\mu, \nu \in \Lambda V$ with $s(\mu) = s(\nu)$, $r(\mu) = r(\nu)$ and $d(\mu) \wedge d(\nu) = m \wedge n = 0$. (If no such μ and ν exist, then Condition (C) is satisfied, [7, Remark 3.2].) Choose open neighborhoods $U_m \subset \Lambda^m$ and $U_n \subset \Lambda^n$ of μ and ν respectively such that $s_m|_{U_m}$ and $s_n|_{U_n}$ are homeomorphisms. Let $V_1 = s_m(U_m) \cap s_n(U_n) \subset V$. Use (B) to choose $\lambda \in V_1\Lambda$ with $d(\lambda) \geq m \vee n = m + n$ that satisfies $(\star)$ for m and n. Take α to be the unique element in $U_m r(\lambda)$ and β the unique element in $U_n r(\lambda)$. Consider

$$\begin{aligned}\alpha\lambda\,(m+n, m+n+d(\lambda) - (m \vee n)) &= \lambda\,(n, n + d(\lambda) - (m \vee n)) \\ &\neq \lambda\,(m, m + d(\lambda) - (m \vee n)) \\ &= \beta\lambda\,(m+n, m+n+d(\lambda) - (m \vee n))\,.\end{aligned}$$

Since

$$\alpha\lambda\,(m+n, m+n+d(\lambda) - (m \vee n)) \neq \beta\lambda\,(m+n, m+n+d(\lambda) - (m \vee n))\,,$$

$\alpha\lambda$ and $\beta\lambda$ can have no common extensions.

(C) $\implies$ (B). Fix an open set $V \subset \Lambda^0$ and a pair $m \neq n \in \mathbb{N}^k$. Let U be the open set $r^{-1}_{m\vee n}(V) \subset \Lambda^{m\vee n}$, choose an open subset $U' \subset U$ so that $s|_{U'}$ is a homeomorphism, and let V' be the open set $s(U')$. By (C) choose $v \in V'$ and $\alpha, \beta \in \Lambda v$ such that there exists a $\tau_0 \in v\Lambda$ with $\mathrm{MCE}(\alpha\tau_0, \beta\tau_0) = \emptyset$. Since s is a homeomorphism onto V', there is a unique $\lambda_0 \in U'$ with $s(\lambda_0) = v$ so $\lambda_0(m \vee n - d(\alpha), m \vee n) = \alpha$ and $\lambda_0(m \vee n - d(\beta), m \vee n) = \beta$. We let $\mu = \lambda_0(m, m \vee n)$ and $\nu = \lambda_0(n, m \vee n)$ and by (C), choose $\tau_0 \in s(\lambda_0)\Lambda$ such that $\mathrm{MCE}(\mu\tau_0, \nu\tau_0) = \emptyset$. Let $\tau_1 \in s(\tau_0)\Lambda^{m\vee n}$ and $\lambda = \lambda_0\tau_0\tau_1$. So,

$$\begin{aligned} d(\lambda) &= d(\lambda_0) + d(\tau_0) + d(\tau_1) \\ &= (m \vee n) + d(\tau_0) + (m \vee n) \\ d(\lambda) - (m \vee n) &= d(\tau_0) + (m \vee n) \end{aligned}$$

Then

$$\begin{aligned} &\lambda(m, m + d(\lambda) - (m \vee n)) \\ &\quad = \lambda(m, (m \vee n) + d(\tau_0) + m) \\ &\quad = \lambda(m, m \vee n)\lambda(m \vee n, m \vee n + d(\tau_0))\lambda(m \vee n + d(\tau_0), m \vee n + d(\tau_0) + m) \\ &\quad = \mu\tau_0\tau_1(0, m) \end{aligned}$$

and similarly $\lambda(n, n + d(\lambda) - (m \vee n)) = \nu\tau_0\tau_1(0, n)$. If $\mu\tau_0\tau_1(0, m) = \nu\tau_0\tau_1(0, n)$, then $\mathrm{MCE}(\mu\tau_0, \nu\tau_0) \neq \emptyset$. Thus, λ must satisfy $(\star)$.

□

While Conditions (B) and (C) may seem complicated or arbitrary, think of their implications in directed graphs. If a directed graph E satisfies Condition (B), then for any two natural numbers $m \neq n$ we can find a path in E such that the m^{th} and n^{th} edges are different. There is also an enlightening diagram of Condition (B) in the discrete case in Appendix A of [8]. Consider two paths μ and ν in a directed graph E where $s(\mu) = s(\nu)$, $r(\mu) = r(\nu)$ and $d(\mu) \wedge d(\nu) = 0$ as described above. This implies that one path, say μ is just a vertex and ν must be a loop based at that vertex. Condition (C) then says that the loop μ must have an entry.

4. Examples

To demonstrate the importance of these formulations of aperiodicity we give an example of a topological k-graph in which Condition (C) is straightforward to verify where as Condition (A) would be quite difficult. First we define what we call a *twisted product topological k-graph.* This twisting construction takes a discrete k-graph and puts a copy of an appropriate topological space X at each vertex, and twists the edges according to a functor τ.

Proposition 4.1. *Let (Λ, d) be a finitely aligned k-graph with no sources, X be a second countable, locally compact, Hausdorff space, and*

$$\tau : \Lambda \to \{\tau_\lambda : X \to X \,|\, \tau_\lambda \text{ is a local homeomorphism}\}$$

a continuous functor. Then the pair $\left(\Lambda \times_\tau X, \tilde{d}\right)$, with object and morphism sets

$$\mathrm{Obj}(\Lambda \times_\tau X) := \mathrm{Obj}(\Lambda) \times X \text{ and } \mathrm{Mor}(\Lambda \times_\tau X) := \mathrm{Mor}(\Lambda) \times X,$$

range and source maps

$$r(\lambda, x) := (r(\lambda), \tau_\lambda(x)) \text{ and } s(\lambda, x) := (s(\lambda), x)\,,$$

and composition

$$(\lambda, \tau_\mu(x)) \circ (\mu, x) = (\lambda\mu, x),$$

whenever $s(\lambda) = r(\mu)$ in (Λ, d), and degree functor

$$\tilde{d}(\lambda, x) = d(\lambda)$$

is a topological k-graph.

The proof that $\left(\Lambda \times_\tau X, \tilde{d}\right)$ satisfies all the requirements of a topological k-graph is straightforward and the details unenlightening, so we omit the proof here.

Consider the discrete 2-graph of Evans and Sims whose 1-skeleton is pictured in Figure 3.



FIGURE 3. The 1-skeleton of a discrete k-graph Λ.

with factorization rules given by

$$\alpha_i^n \beta_j^{n+1} = \beta_{\xi_n(i,j)}^n \alpha_{\zeta_n(i,j)}^{n+1}$$

where $(i, j) \mapsto (\xi_n(i, j), \zeta_n(i, j))$ is a permutation of $\mathbb{Z}_n \times \mathbb{Z}_{n+1}$.

For an example of a twisted product we take the 1-skeleton of Figure 3, with ξ_n and ζ_n being “plus 1” in the group $\mathbb{Z}/n\mathbb{Z}$, the topological space $\mathbb{T}$ with the usual topology, and the functor τ given by $\tau_{\alpha_i^n}(z) = \tau_{\beta_j^n}(z) := z^n$.

First, we check if the twisted topological k-graph above satisfies Condition (C). Similar to the proof of (B) $\implies$ (C), we fix an open set $V \subset \Lambda^0$ and paths $\mu, \nu \in V\Lambda$ such that $r(\mu) = r(\nu)$, $s(\mu) = s(\nu)$ and $d(\mu) \wedge d(\nu) = 0$. We can see that μ must have the form

$$\mu = \left(\alpha^m_{i_0}, z^k\right) \dots \left(\alpha^{m+n-2}_{i_{n-2}}, \left(z^{m+n}\right)^{m+n-1}\right)\left(\alpha^{m+n-1}_{i_{n-1}}, z^{m+n}\right)\left(\alpha^{m+n}_{i_n}, z\right)$$

where k is given by $k = \prod_{i=m+1}^{m+n} i$. Similarly,

$$\nu = \left(\beta^m_{j_0}, z^k\right) \dots \left(\beta^{m+n}_{j_n}, z\right)$$

We consider the path $\lambda = \left(\beta^{m+n+1}_{j_0-n}, z^{1/m+n+1}\right)$. Notice the path $\nu\lambda$ is built of all "β" edges. We follow the factorization rules to rewrite the path $\mu\lambda$ such that the first edge is a "β" edge, and then compare to $\nu\lambda$.

$$\begin{aligned}\mu\lambda &= \left(\alpha^m_{i_0}, z^k\right)\left(\alpha^{m+1}_{i_1}, z^{k/m+n}\right) \dots \left(\alpha^{m+n}_{i_n}, z\right)\left(\beta^{m+n+1}_{j_0-n}, z^{1/m+n+1}\right) \\ &= \left(\beta^m_{j_0+1}, z^k\right)\left(\alpha^{m+1}_{i_0+1}, z^{k/m+n}\right) \dots \left(\alpha^{m+n}_{i_n+1}, z^{1/m+n+1}\right)\end{aligned}$$

If $m \neq 1$, $\left(\beta^m_{j_0}, z^k\right) \neq \left(\beta^m_{j_0+1}, z^k\right)$ and the paths $\mu\lambda$ and $\nu\lambda$ will have no common extensions. If $m = 1$, then we'd amend a similar path of shape $(0, 2)$ so that $\mu\lambda$ and $\nu\lambda$ would be guaranteed to differ in the second β-edge by a similar calculation. Thus, $\mathrm{MCE}(\mu\lambda, \nu\lambda) = \emptyset$ and $\Lambda \times_\tau \mathbb{T}$ is aperiodic.

In contrast, if we were to check for aperiodicity using Condition (A) we would need to consider infinite paths of the form

$$x = \left(\alpha^i_0, z\right)\left(\beta^{i+1}_0, z^{1/i+1}\right)\left(\alpha^{i+2}_0, z^{1/(i+1)(i+2)}\right)\left(\beta^{i+3}_0, z^{1/(i+1)(i+2)(i+3)}\right)\dots$$

and investigate $\sigma^m x$ for any $m \in \mathbb{N}^k$. A precise calculation requires a variety of formulas for $\sigma^m x$ to cover different cases and then checking the various factorizations of the result for periodic behavior. It is at least three long and some what complicated calculations.

This demonstrates how much easier it is to check aperiodicity through Condition (C) rather than (A) even in this fairly straightforward example. Imagine starting with a discrete Λ with a more complicated permutations, constructing a twisted product with a space more complicated than $\mathbb{T}$, or a twisting functor that is more varied than the τ above. It can only get more difficult.

References

1. Joachim Cuntz and Wolfgang Krieger, *A class of C^*-algebras and topological Markov chains*, Invent. Math. **56** (1980), no. 3, 251–268. MR MR561974 (82f:46073a)
2. Masatoshi Enomoto and Yasuo Watatani, *A graph theory for C^*-algebras*, Math. Japon. **25** (1980), no. 4, 435–442. MR MR594544 (83d:46069a)
3. Cynthia Farthing, Nura Patani, and Paulette Willis, *The crossed-product structure of C^*-algebras arising from topological dynamical systems*, to appear.

4. Takeshi Katsura, *A class of C^*-algebras generalizing both graph algebras and homeomorphism C^*-algebras. I. Fundamental results*, Trans. Amer. Math. Soc. **356** (2004), no. 11, 4287–4322 (electronic).
5. Alex Kumjian and David Pask, *Higher rank graph C^*-algebras*, New York J. Math. **6** (2000), 1–20 (electronic). MR MR1745529 (2001b:46102)
6. Alex Kumjian, David Pask, Iain Raeburn, and Jean Renault, *Graphs, groupoids, and Cuntz-Krieger algebras*, J. Funct. Anal. **144** (1997), no. 2, 505–541. MR MR1432596 (98g:46083)
7. Peter Lewin and Aidan Sims, *Aperiodicity and cofinality for finitely aligned higher-rank graphs*, Math. Proc. Cambridge Philos. Soc. **149** (2010), no. 2, 333–350. MR 2670219
8. David I. Robertson and Aidan Sims, *Simplicity of C^*-algebras associated to higher-rank graphs*, Bull. Lond. Math. Soc. **39** (2007), no. 2, 337–344. MR MR2323468 (2008g:46099)
9. Sarah Wright, *Aperiodicity in topological k-graphs*, Ph.D. thesis, May 2010.
10. Trent Yeend, *Topological higher-rank graphs and the C^*-algebras of topological 1-graphs*, Operator theory, operator algebras, and applications, Contemp. Math., vol. 414, Amer. Math. Soc., Providence, RI, 2006, pp. 231–244. MR MR2277214 (2007j:46100)
11. ______, *Groupoid models for the C^*-algebras of topological higher-rank graphs*, J. Operator Theory **57** (2007), no. 1, 95–120. MR MR2301938 (2008f:46074)
12. ______, *Topological higher-rank graphs, their groupoids and operator algebras*, Ph.D. thesis, March 2005.

College of the Holy Cross
E-mail address: `swright@holycross.edu`